


































HEEL MEDICATIONS

Allergies

Reference Book



Main Heel products for the following symptoms and conditions	Navigate by key products									
	Lufeel® nasal spray	Lufeel® tabs.	Cutis compositum	Oculoheel® (drops)	Hepeel®	Mucosa comp.	Engystol®	Coenzyme composition	Lymphomyosot®	
Main product  Supportive product 										
Sneezing										
Rhinorrhea										
Nasal & throat/mucous membrane itching										
Nasal obstruction										
Nasal & sinus congestion										
Nasal polyps										
Skin symptoms: itching, papules, hives, localized swelling*										
Ocular symptoms: burning, itching, aching, redness, lacrimation										
Targeting the pathophysiology in addition to above treatment recommendations										
Support of liver in removing waste products triggering allergy, e.g. histamine										
Regulation and support of mucosal membrane functions										
Restoring balance in the immune system										
Support of cellular respiration in chronic disease										
Prophylaxis in seasonal allergies (Start 12 weeks before allergy season)										

*In severe eczema, immunomodulation with Engystol and Cutis compositum should be done for at least 4 weeks before the detoxification with Lymphomyosot is started

Product	Indications	
Luffeel®/Luffa comp.	<ul style="list-style-type: none"> Allergic rhinitis 	5.4
Cutis compositum	<ul style="list-style-type: none"> Dermatitis Allergic dermatitis Seborrheic dermatitis Neurodermatitis 	5.7
Oculoheel®	<ul style="list-style-type: none"> Conjunctivitis Eye irritation Over-strained eyes 	5.8
Hepeel®	<ul style="list-style-type: none"> Improves metabolic function of the liver Supports detoxification 	5.9
Mucosa compositum	<ul style="list-style-type: none"> Regulation and support of the mucosal membrane functions 	5.10
Engystol®	<ul style="list-style-type: none"> Supports immune system 	5.11
Coenzyme compositum	<ul style="list-style-type: none"> Stimulation of blocked intracellular enzymatic systems 	5.12
Lymphomyosot®	<ul style="list-style-type: none"> Lymph flow disorders Recurrent infections Drainage and Detoxification 	5.13



Fight symptoms of allergies and hay fever

- Allergic rhinitis (treatment and prophylaxis)



Dosage Recommendations



Nasal spray

In general: 1-2 shots into each nostril 3-5 times daily
 Children under 6 years: 1 shot 3-4 times daily

For further information, see package insert.

Shield yourself from allergies and hay fever

- Allergic rhinitis (treatment and prophylaxis)



Dosage Recommendations




Tablets

In general: 1 tablet 4 times daily
 In acute disorders: 1 tablet every 15 minutes, over a period lasting up to 2 hours

For further information, see package insert.

Clinical Evidence

- In a randomized double blind equivalence trial with 146 patients, the efficacy and tolerance of Luffa comp.-Heel nasal spray was compared with Cromolyn sodium spray in the treatment of seasonal allergic rhinitis (hay fever).



Reference:

Weiser M., Gegenheimer L.H., Klein P. Seasonal allergic rhinitis: A homeopathic nasal spray as effective as cromoglicic acid. *Frosch Komplementärmed/ Research in Complementary Medicine*. 1999;6:142-148

- Luffa comp.-Heel showed a comparable onset of action
- Both treatments had their main therapeutic effect in the first week
- No adverse systemic effects (e.g. drowsiness, fatigue) and very rare local adverse effects (e.g. smarting of the nostrils)


Patient Profiles

- **Kate | 19-year-old**

Single

Medical history: atopic dermatitis as a child confirmed with raised specific IgE antibodies to pollen, dander and dust mites.

Present Disorders: allergic rhinoconjunctivitis with itching, redness, aching and lacrimation of the eyes when exposed to the above allergens.



- **Tina | 26-year-old**

Married, 1 child

Job: school teacher

Medical history: positive skin prick tests to pollen and certain grass types.

Present disorders: sneezing attacks, rhinorrhea, itchy palate and ears followed by nasal congestion that interferes with her quality of life during the allergy season. The patient would like a treatment that can possibly assist in preventing the symptoms.



- **Roger | 32-year-old**

Single

Occupation: Artist

Family history: mother suffers with atopic dermatitis when stressed and sister suffers with seasonal hay fever

Medical history: hay fever since a teenager and chronic sinusitis sufferer.

Present disorders: episodes of sneezing and rhinorrhea with chronic nasal and sinus congestion with malaise and drowsiness.







Comprehensive treatment of skin disorders

- Dermatitis
- Allergic dermatitis
- Seborrheic dermatitis
- Neurodermatitis



Dosage Recommendations

		ACUTE DOSAGE	STANDARD DOSAGE
 Tablets	Adults (and children > 12 years)	1 tablet every 1/2 to 1 hr. for the first day up to 12x then continue with 1 tablet 3x/dail	1 tablet 3x daily
	Children 6-11 years	1 tablet every 1/2 to 1 hr. for the first day	1 tablet 2x daily
	Children 2-5 years	1 tablet every 1/2 to 1 hr. for the first day	1 tablet 1-2x daily
	Children 0-2 years	1 tablet every 1/2 to 1 hr. for the first day	1 tablet 1x daily
 Injection solution i.m., s.c., i.d., i.v.	Adults (and children > 12 years)	1 ampoule daily	1 ampoule 1-3x weekly
	Children 6-11 years	1 ampoule daily	2/3 ampoule 1-3x weekly
	Children 2-5 years	1 ampoule daily	1/2 ampoule 1-3x weekly
For further information, see package insert.			


Oculoheel®

Calm an inflamed allergic eye

- Conjunctivitis
- Eye irritation
- Over-strained eyes



Dosage Recommendations

		ACUTE DOSAGE	STANDARD DOSAGE
 Eye drops	Adults (and children > 12 years)	–	1 drop to drip into the affected eye, 3x daily
	Children 6-11 years	–	1 drop 3 times daily
	Children 2-5 years	–	1 drop 1-2x daily
	Children 0-2 years	–	1 drop 1x daily
For further information, see package insert.			





Support the removal of waste products that trigger allergies

- Improves metabolic function of the liver
- Supports detoxification



Dosage Recommendations

		ACUTE DOSAGE	STANDARD DOSAGE
 <p>Tablets</p>	Adults (and children > 12 years)	1 tablet every 1/2-1 hour, up to 12x daily	1 tablet dissolved in the mouth 3x daily
	Children 6-11 years	1 tablet every 1-2 hours, up to 8x daily	1 tablet 2x daily
	Children 2-5 years	1 tablet every 1-2 hours, up to 6x daily	1 tablet 1-2x daily
	Children 0-2 years	1 tablet every 1-2 hours, up to 4x daily	1 tablet 1x daily
 <p>Injection solution i.m., s.c., i.d., i.v.</p>	Adults (and children > 12 years)	1 ampoule daily for 3 days, then continue with standard dosage	1 ampoule 1-3x weekly
	Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly
	Children 2-5 years	1/2 ampoule daily	1/2 ampoule 1-3x weekly
For further information, see package insert.			



Mucosa compositum

For healthy functioning
of the mucosal membranes

- Regulation and support of the mucosal membrane functions, including the gastrointestinal tract



Dosage Recommendations

		ACUTE DOSAGE	STANDARD DOSAGE
 Tablets	Adults (and children > 12 years)	1 tablet every 1/2-1 hour, up to 12x daily	1 tablet 3x daily
	Children 6-11 years	1 tablet every 1-2 hours, up to 8x daily	1 tablet 2x daily
	Children 2-5 years	1 tablet every 1-2 hours, up to 6x daily	1 tablet 1-2x daily
	Children 0-2 years	1 tablet every 1-2 hours, up to 4x daily	1 tablet 1x daily
 Injection solution i.m., s.c., i.d., i.v.	Adults (and children > 12 years)	1 ampoule daily for 3 days, then continue with standard dosage	1 ampoule 1-3x weekly
	Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly
	Children 2-5 years	1/2 ampoule daily	1/2 ampoule 1-3x weekly
For further information, see package insert.			





**Support your immune system
in fighting off viral infections,
including colds and flu**

- Common colds
- Mild viral infections
- Flu-like illnesses



Dosage Recommendations

		ACUTE DOSAGE	CHRONIC DOSAGE
 Tablets	Adults (and children > 12 years)	1 tablet every 1/2-1 hour, up to 12x daily	1 tablet 3x daily
	Children 6-11 years	1 tablet every 1-2 hours, up to 8x daily	1 tablet 2x daily
	Children 2-5 years	1 tablet every 1-2 hours, up to 6x daily	1 tablet 1-2x daily
	Children 0-2 years	1 tablet every 1-2 hours, up to 4x daily	1 tablet 1x daily
 Injection solution i.m., s.c., i.d., i.v.	Adults (and children > 12 years)	1 ampoule daily	1 ampoule 1-3x weekly
	Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly
	Children 2-5 years	1/2 ampoule daily	1/2 ampoule 1-3x weekly
For further information, see package insert.			





A multi target cell tuner

- Stimulation of blocked intracellular enzymatic systems



Dosage Recommendations

		ACUTE DOSAGE	STANDARD DOSAGE
 <p>Tablets</p>	Adults (and children > 12 years)	1 tablet every 1/2-1 hour, up to 12x daily	1 tablet 3x daily
	Children 6-11 years	1 tablet every 1-2 hours, up to 8x daily	1 tablet 2x daily
	Children 2-5 years	1 tablet every 1-2 hours, up to 6x daily	1 tablet 1-2x daily
	Children 0-2 years	1 tablet every 1-2 hours, up to 4x daily	1 tablet 1x daily
 <p>Injection solution i.m., s.c., i.d., i.v.</p>	Adults (and children > 12 years)	1 ampoule daily	1 ampoule 1-3x weekly
	Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly
	Children 2-5 years	1/2 ampoule daily	1/2 ampoule 1-3x weekly
For further information, see package insert.			





Restore the lymphatic flow for a faster recovery

- Lymph flow disorders
- Recurrent infections
- Drainage and Detoxification



Dosage Recommendations

		ACUTE DOSAGE	STANDARD DOSAGE
 Tablets	Adults (and children > 12 years)	1 tablet every 1/2-1 hour, up to 12x daily	1 tablet 3x daily
	Children 6-11 years	1 tablet every 1-2 hours, up to 8x daily	1 tablet 2x daily
	Children 2-5 years	1 tablet every 1-2 hours, up to 6x daily	1 tablet 1-2x daily
	Children 0-2 years	1 tablet every 1-2 hours, up to 4x daily	1 tablet 1x daily
 Injection solution i.m., s.c., i.d., i.v.	Adults (and children > 12 years)	1 ampoule daily	1 ampoule 1-3x weekly
	Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly
	Children 2-5 years	1/2 ampoule daily	1/2 ampoule 1-3x weekly
For further information, see package insert.			

Product Information

Coenzyme comp. • Tablets • Injection solution

Compositions: Tablets: 1 tablet = 301.5 mg containing: Active ingredients: Acidum cis-aconiticum D8 1.0 mg, Acidum ascorbicum D6 1.0 mg, Acidum citricum D8 1.0 mg, Acidum fumaricum D8 1.0 mg, Acidum alpha-ketoglutaricum D8 1.0 mg, Acidum malicum D8 1.0 mg, Acidum succinicum D8 1.0 mg, Adenosinum triphosphoricum D10 1.0 mg, Beta vulgaris rubra D6 1.0 mg, Coenzym A D8 1.0 mg, Cysteinum D6 1.0 mg, Nadidum D8 1.0 mg, Natrium pyruvicum D8 1.0 mg, Natrium riboflavinum phosphoricum D6 1.0 mg, Nicotinamidum D6 1.0 mg, Pulsatilla pratensis D6 1.0 mg, Pyridoxinum hydrochloricum D6 1.0 mg, Sulfur D10 1.0 mg, Thiaminum hydrochloricum D6 1.0 mg, Acidum thiocticum D6 1.0 mg, Barium oxalsuccinicum D10 1.0 mg, Cerium oxalicum D8 1.0 mg, Hepar sulfuris D10 1.0 mg, Magnesium oroticum dihydricum D6 1.0 mg, Manganum phosphoricum D6 1.0 mg, Natrium diethylxalacetatum D6 1.0 mg. Excipients: Lactose monohydrate 293.0 mg, Magnesium stearate 1.5 mg. **Injection solution:** 2.2 g containing: Acidum ascorbicum D6 22.0 mg, Acidum alpha-ketoglutaricum D8 22.0 mg, Acidum cis-aconiticum D8 22.0 mg, Acidum citricum D8 22.0 mg, Acidum fumaricum D8 22.0 mg, Acidum malicum D8 22.0 mg, Acidum succinicum D8 22.0 mg, Adenosinum triphosphoricum D10 22.0 mg, Barium oxalsuccinicum D10 22.0 mg, Beta vulgaris rubra D4 22.0 mg, Coenzym A D8 22.0 mg, Cysteinum D6 22.0 mg, Hepar sulfuris D10 22.0 mg, Nadidum D8 22.0 mg, Natrium pyruvicum D8 22.0 mg, Natrium riboflavinum phosphoricum D6 22.0 mg, Nicotinamidum D6 22.0 mg, Pulsatilla pratensis D6 22.0 mg, Pyridoxinum hydrochloricum D6 22.0 mg, Sulfur D10 22.0 mg, Thiaminum hydrochloricum D6 22.0 mg, Acidum thiocticum D6 22.0 mg, Cerium oxalicum D8 22.0 mg, Magnesium oroticum dihydricum D6 22.0 mg, Manganum phosphoricum D6 22.0 mg, Natrium diethylxalacetatum D6 22.0 mg. Excipients: Sodium chloride 19.4 mg, water for injections 1628.0 mg.

Indications: Tablets, injection solution: Stimulation of blocked intracellular respiratory enzymatic systems in degenerative diseases.

Contraindications: Tablets, injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Injection solution:** None.

Side effects: Tablets: None have been reported. **Injection solution:** Allergic (hypersensitivity) reactions (e.g. skin allergies, redness/swelling at the injection site) may occur in very rare cases (i.e. affects less than 1 in 10,000 users).

Interactions with other medication: Tablets, injection solution: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. Acute or Initial Dosage: Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. Method of Administration: Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Injection solution:** Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ½ of an ampoule 1 to 3x weekly. Acute or Initial Dosage: Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ½ of an ampoule daily, and then continue with standard dosage. Method of Administration: Coenzyme comp., Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Tablets, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (41946): Packs containing 50 and 250 tablets. **Injection solution (8511):** Packs containing 5, 10, 50 and 100 ampoules of 2.2 ml each.

Cutis comp. • Tablets • Injection solution

Compositions: Tablets: 1 tablet = 301.5 mg containing: Active ingredients: Acidum formicicum D198 1.0 mg, Acidum fumaricum D10 1.0 mg, Acidum phosphoricum D6 1.0 mg, Aesculus hippocastanum D6 1.0 mg, Ammonium bituminosulfonicum D28 1.0 mg, Arctium D6 1.0 mg, Cortisonum aceticum D28 1.0 mg, Cutis suis D8 1.0 mg, Funiculus umbilicalis suis D10 1.0 mg, Galium aparine D6 1.0 mg, Glandula suprarenalis suis D10 1.0 mg, Hepar suis D10 1.0 mg, Ledum palustre D4 1.0 mg, Placenta totalis suis D10 1.0 mg, Splen suis D10 1.0 mg, Strychnos ignatii D6 1.0 mg, Sulfur D10 1.0 mg, Thuja occidentalis D8 1.0 mg, Urtica urens D4 1.0 mg, Calcium fluoratum D13 1.0 mg, Mercurius solubilis Hahnemanni D13 1.0 mg, Selenium D10 1.0 mg. Excipients: Lactose monohydrate 297.0 mg, Magnesium stearate 1.5 mg. **Injection solution:** 2.2 g containing: Active ingredients: Acidum alpha-ketoglutaricum D10 22.0 mg, Acidum formicicum D198 22.0 mg, Acidum fumaricum D10 22.0 mg, Acidum phosphoricum D6 22.0 mg, Aesculus hippocastanum D6 22.0 mg, Ammonium bituminosulfonicum D28 22.0 mg, Arctium D6 22.0 mg, Calcium fluoratum D13 22.0 mg, Cortisonum aceticum D28 22.0 mg, Cutis suis D8 22.0 mg, Funiculus umbilicalis suis D10 22.0 mg, Galium aparine D6 22.0 mg, Glandula suprarenalis suis D10 22.0 mg, Hepar suis D10 22.0 mg, Ledum palustre D4 22.0 mg, Mercurius solubilis Hahnemanni D13 22.0 mg, Natrium diethylxalacetatum D10 22.0 mg, Placenta totalis suis D10 22.0 mg, Selenium D10 22.0 mg, Splen suis D10 22.0 mg, Strychnos ignatii D6 22.0 mg, Sulfur D10 22.0 mg, Thallium sulfuricum D13 22.0 mg, Thuja occidentalis D8 22.0 mg, Urtica urens D4 22.0 mg. Excipients: Sodium chloride 19.4 mg, water for injections 1650.0 mg.

Indications: Tablets, injection solution: Supportive treatment of chronic skin conditions.

Contraindications: Tablets, injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Injection solution:** None.

Side effects: Tablets, injection solution: Like all medicinal products, homeopathic medicines can cause side effects in isolated cases, such as transient allergic reactions. The frequency of these effects is not known.

Interactions with other medication: Tablets, injection solution: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. Method of administration: Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Injection solution:** Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ½ of an ampoule 1 to 3x weekly. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ½ of an ampoule daily, and then continue with standard dosage. Method of administration: Cutis comp., Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Tablets, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (44179): Packs containing 50 and 250 tablets. **Injection solution (9477):** Packs containing 5, 10, 50 and 100 ampoules of 2.2 ml each.

Engystol® • Tablets • Injection solution

Compositions: Tablets: 1 tablet = 301.5 mg containing: Active ingredients: Vincetoxicum hirundinaria D6, Vincetoxicum hirundinaria D10, Vincetoxicum hirundinaria D30 – 75 mg each; Sulfur D4, Sulfur D10 – 37.5 mg each. Excipients: Contains lactose; Magnesium stearate 1.5 mg. **Injection solution:** 1.1 g containing: Active ingredients: Vincetoxicum hirundinaria D6, Vincetoxicum hirundinaria D10, Vincetoxicum hirundinaria D30 – 6.6 mg each; Sulfur D4, Sulfur D10 – 3.3 mg each. Excipients: Sodium chloride 8.78 mg, water for injections 1073.60 mg

Indications: Tablets, injection solution: Supports the immune system to reduce severity and duration of symptoms in viral infections, particularly in colds and influenza-like illnesses.

Contraindications: Tablets, injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Injection solution:** None.

Side effects: Tablets, injection solution: Allergic (hypersensitivity) reactions (e.g. skin allergies, redness/swelling at the injection site) may occur in very rare cases (i.e. affects less than 1 in 10,000 users).

Interaction with other medicaments: Tablets, injection solution: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs. up to 8x daily, and then continue with standard dosage. Method of administration: Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Injection solution:** Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅓ of an ampoule 1 to 3x weekly. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅓ of an ampoule daily, and then continue with standard dosage. Method of administration: Injection solution may be administered by the s.c., i.d., i.m., or i.v. route.

Overdose: Tablets, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (9830): Packs containing 50 and 250 tablets. **Injection solution (8546):** Packs containing 5, 10, 50 and 100 ampoules of 1.1 ml.

Hepeel® • Tablets • Injection solution

Compositions: Tablets: 1 tablet (301.5 mg) contains: Active ingredients: Chelidonium majus D6 30.0 mg, Cinchona pubescens D3 30.0 mg, Citrullus colocynthis D6 90.0 mg, Lycopodium clavatum D3 30.0 mg, Myristica fragrans D4 30.0 mg, Phosphorus D6 15.0 mg, Silybum marianum D2 15.0 mg, Veratrum album D6 60.0 mg. Excipients: Magnesium stearate 1.5 mg. Contains lactose. **Injection solution:** 1 ampoule (1.1 g) containing: Active ingredients: Chelidonium majus D6 1.1 mg, Cinchona pubescens D2 1.1 mg, Citrullus colocynthis D5 3.3 mg, Lycopodium clavatum D2 1.1 mg, Myristica fragrans D3 1.1 mg, Phosphorus D5 0.55 mg, Silybum marianum D1 0.55 mg, Veratrum album D5 2.2 mg. Excipients: Sodium chloride 8.87–9.58 mg, water for injections 1089.00 mg.

Indications: Tablets, injection solution: Functional disorders of the hepatobiliary system, e.g. in hepatitis; hepatoprotectant.

Contraindications: Tablets, injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. In case of a history of liver diseases, appropriate standards of care should be followed. **Injection solution:** In case of a history of liver diseases, appropriate standards of care should be followed.

Side effects: Tablets, injection solution: Like all medicinal products, homeopathic medicines can cause side effects in isolated cases, such as transient allergic reactions. The frequency of these effects is not known.

Interactions with other medication: Tablets, injection solution: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. Method of administration: Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Injection solution:** Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅓ of an ampoule 1 to 3x weekly. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅓ of an ampoule daily, and then continue with standard dosage. Method of administration: Hepeel, Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Tablets, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (47787): Packs containing 50 and 250 tablets. **Injection solution (46791):** Packs containing 5, 10, 50 and 100 ampoules of 1.1 ml each.

Luffeel® • Tablets • Nasal spray

Compositions: Tablets: 301.5 mg contains: Active ingredients: Arsenum iodatum D8 25 mg, Lobelia inflata D6 25 mg, Luffa operculata D12 25 mg, Aralia racemosa D1 25 mg. Excipients: Lactose monohydrate 200 mg, Magnesium stearate 1.5 mg. **Nasal spray:** 100 g contains: Active ingredients: Histaminum D12 5 g, Histaminum D30 5 g, Histaminum D200 5 g, Luffa operculata D4 10 g, Luffa operculata D12 10 g, Luffa operculata D30 10 g, Sulfur D12 5 g, Sulfur D30 5 g, Sulfur D200 5 g, Thryallis glauca D4 10 g, Thryallis glauca D12 10 g, Thryallis glauca D30 10 g. Excipients: Benzalkonium chloride solution 0.02 g, Disodium phosphate dihydrate 0.02 g, Sodium chloride 0.72 g, Sodium dihydrogen phosphate dihydrate 0.0628 g, Water purified 9.1772 g.

Indications: Tablets, Nasal spray: Allergic rhinitis, particularly hay fever. May also be used to support the preventative (prophylactic) management of these conditions. **Nasal spray:** Allergic rhinitis, particularly hay fever.

Contraindications: Tablets, Nasal spray: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Nasal spray:** Luffeel, Nasal spray contains benzalkonium chloride. In some patients this may cause irritation of the nasal mucosa. This occurs very rarely (i.e. affects less than 1 in 10.00 users). In very rare cases bronchospasm may occur in predisposed asthmatic patients due to benzalkonium chloride (i.e. affects less than 1 in 10.00 users).

Side effects: Tablets: Allergic (hypersensitivity) skin reactions may occur in very rare cases (i.e. affects less than 1 in 10.000 users). **Nasal spray:** Local allergic (hypersensitivity) reactions or irritation of the nasal mucosa may occur in very rare cases (i.e. affects less than 1 in 10.000 users).

Interactions with other medication: Tablets, Nasal spray: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, Nasal spray: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during the pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, Nasal spray: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard Dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. Acute or Initial Dosage: Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. Preventative (Prophylactic) Dosage: For patients in whom allergy symptoms recur annually, start with the standard dosage one month before the allergy season and continue until two weeks after the end of the critical season. Method of administration: Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add a small amount of water. This medicine should be taken away from meals. **Nasal spray:** Standard Dosage: Adults (and children 12 yrs. and older): 1–2 sprays into each nostril 3–5x daily. Pediatric: 2–5 yrs.: 1 spray into each nostril 3–5x daily. 6–11 yrs.: 1 spray into each nostril 3–5x daily. Acute or Initial Dosage: Adults (and children 12 yrs. and older): 1–2 sprays into each nostril every ½ to 1 hr. for the first day up to 8x, and then continue with standard dosage. Pediatric: 2–5 yrs.: 1 spray into each nostril every ½ to 1 hr. for the first day up to 5x, and then continue with standard dosage. 6–11 yrs.: 1 spray into each nostril every ½ to 1 hr. for the first day up to 8x, and then continue with standard dosage. Method of administration: For each application, spray the recommended dosage into each nostril. For hygienic reasons it is not recommended to share the use of the same product with others.

Overdose: Tablets, Nasal spray: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (9831): Packs containing 50 and 250 tablets. **Nasal spray (9864):** Atomiser without propellant, 20 ml.

Lymphomyosot® • Tablets • Oral drops • Injection solution

Compositions: Tablets: 1 tablet = 301.5 mg containing: Active ingredients: Araneus diadematus D6 15.0 mg, Calcium phosphoricum D12 15.0 mg, Equisetum hiemale D4 15.0 mg, Ferrum iodatum D12 30.0 mg, Fumaria officinalis D4 15.0 mg, Gentiana lutea D5 15.0 mg, Geranium robertianum D4 30.0 mg, Juglans regia ssp. regia D3 15.0 mg, Levothyroxinum D12 15.0 mg, Myosotis arvensis D3 15.0 mg, Nasturtium officinale D4 30.0 mg, Natrium sulfuricum D4 15.0 mg, Pinus sylvestris D4 15.0 mg, Scrophularia nodosa D3 15.0 mg, Smilax D6 15.0 mg, Teucrium scorodonia D3 15.0 mg, Veronica officinalis D3 15.0 mg. Contains lactose. **Oral drops:** 100 g containing: Active ingredients: Araneus diadematus D6 5.0 g, Calcium phosphoricum D12 5.0 g, Equisetum hiemale D4 5.0 g, Ferrum iodatum D12 10.0 g, Fumaria officinalis D4 5.0 g, Gentiana lutea D5 5.0 g, Geranium robertianum D4 10.0 g, Levothyroxinum D12 5.0 g, Myosotis arvensis D3 5.0 g, Nasturtium officinale D4 10.0 g, Natrium sulfuricum D4 5.0 g, Pinus sylvestris D4 5.0 g, Scrophularia nodosa D3 5.0 g, Smilax D6 5.0 g, Teucrium scorodonia D3 5.0 g, Veronica officinalis D3 5.0 g, Excipients: Water, purified 5 g. Contains 35 vol.-% alcohol. **Injection solution:** 1.1 g containing: Active ingredients: Araneus diadematus D6 0.55 mg, Calcium phosphoricum D12 0.55 mg, Equisetum hiemale D4 0.55 mg, Ferrum iodatum D12 1.10 mg, Fumaria officinalis D4 0.55 mg, Gentiana lutea D5 0.55 mg, Geranium robertianum D4 1.10 mg, Levothyroxinum D12 0.55 mg, Myosotis arvensis D3 0.55 mg, Nasturtium officinale D4 1.10 mg, Natrium sulfuricum D4 0.55 mg, Pinus sylvestris D4 0.55 mg, Scrophularia nodosa D3 0.55 mg, Smilax D6 0.55 mg, Teucrium scorodonia D3 0.55 mg, Veronica officinalis D3 0.55 mg. Excipients: Sodium chloride 10.4 mg, water for injections 1089.6 mg.

Indications: Tablets, oral drops, injection solution: Improvement of lymphatic drainage, the non-specific immune defense, and conditions such as benign hypertrophy of lymph nodes, chronic tonsillitis, tonsillar hypertrophy and lymphatic edema.

Contraindications: Tablets, oral drops, injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. Oral drops: This medicinal product contains 35 vol.-% ethanol (alcohol). **Injection solution:** None.

Side effects: Tablets, oral drops, injection solution: Allergic (hypersensitivity) skin reactions may occur in very rare cases (i.e. affects less than 1 in 10,000 users). **Interactions with other medication: Tablets, oral drops, injection solution:** No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, oral drops, injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, oral drops, injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. Method of administration: Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Oral drops:** Standard dosage: Adults (and children 12 yrs. and older): 10 drops 3x daily. Pediatric: Below 2 yrs.: 3 drops 3x daily. 2–5 yrs.: 5 drops 3x daily. 6–11 yrs.: 7 drops 3x daily. Acute or initial dosage: Adults (and children 12 yrs. and older): 10 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 3 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. 2–5 yrs.: 5 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. 6–11 yrs.: 7 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Method of administration: This medicine should be taken away from meals. For children, add drops to a small amount of water. **Injection solution:** Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅓ of an ampoule 1 to 3x weekly. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅓ of an ampoule daily, and then continue with standard dosage. Method of administration: Lymphomyosol, Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Tablets, drops, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: (9832): Packs containing 50, 100 and 250 tablets. **Oral drops (10019):** Packs containing 30 ml and 100 ml. **Injection solution (9487):** Packs containing 5, 10, 50 and 100 ampoules of 1.1 ml each.

Mucosa compositum • Tablets • Injection solution

Compositions: Tablets: 1 tablet = 301.5 mg containing: Active ingredients: Argentum nitricum D6 1.0 mg, Atropa bella-donna D10 1.0 mg, Ceanothus americanus D4 1.0 mg, Hydrastis canadensis D6 1.0 mg, Kalium bichromicum D8 1.0 mg, Kreosotum D12 1.0 mg, Lachesis D10 1.0 mg, Mandragora e radice siccata D10 1.0 mg, Marsdenia cundurango D6 1.0 mg, Momordica balsamina D6 1.0 mg, Mucosa coli suis D8 1.0 mg, Mucosa ductus choledochi suis D8 1.0 mg, Mucosa duodeni suis D8 1.0 mg, Mucosa ilei suis D8 1.0 mg, Mucosa jejuni suis D8 1.0 mg, Mucosa nasalis suis D8 1.0 mg, Mucosa oculi suis D8 1.0 mg, Mucosa oesophagi suis D8 1.0 mg, Mucosa oris suis D8 1.0 mg, Mucosa pulmonis suis D8 1.0 mg, Mucosa pylori suis D8 1.0 mg, Mucosa recti suis D8 1.0 mg, Mucosa vesicae felleae suis D8 1.0 mg, Mucosa vesicae urinae suis D8 1.0 mg, Oxalis acetosella D6 1.0 mg, Pankreas suis D10 1.0 mg, Phosphorus D8 1.0 mg, Psychotria ipecacuanha D8 1.0 mg, Pulsatilla pratensis D6 1.0 mg, Semecarpus anacardium D6 1.0 mg, Strychnos nux-vomica D13 1.0 mg, Sulfur D8 1.0 mg, Ventriculus suis D8 1.0 mg, Veratrum album D6 1.0 mg. Excipients: Lactose monohydrate 300 mg, Magnesium stearate 1.5 mg. **Injection solution:** 2.2 g containing: Active ingredients: Argentum nitricum D6 22.0 mg, Atropa bella-donna D10 22.0 mg, Ceanothus americanus D4 22.0 mg, Hydrastis canadensis D4 22.0 mg, Kalium bichromicum D8 22.0 mg, Kreosotum D10 22.0 mg, Lachesis D10 22.0 mg, Mandragora e radice siccata D10 22.0 mg, Marsdenia cundurango D6 22.0 mg, Momordica balsamina D6 22.0 mg, Mucosa coli suis D8 22.0 mg, Mucosa ductus choledochi suis D8 22.0 mg, Mucosa duodeni suis D8 22.0 mg, Mucosa ilei suis D8 22.0 mg, Mucosa jejuni suis D8 22.0 mg, Mucosa nasalis suis D8 22.0 mg, Mucosa oculi suis D8 22.0 mg, Mucosa oesophagi suis D8 22.0 mg, Mucosa oris suis D8 22.0 mg, Mucosa pulmonis suis D8 22.0 mg, Mucosa pylori suis D8 22.0 mg, Mucosa recti suis D8 22.0 mg, Mucosa vesicae felleae suis D8 22.0 mg, Mucosa vesicae urinae suis D8 22.0 mg, Oxalis acetosella D6 22.0 mg, Pankreas suis D10 22.0 mg, Phosphorus D8 22.0 mg, Psychotria ipecacuanha D8 22.0 mg, Pulsatilla pratensis D6 22.0 mg, Semecarpus anacardium D6 22.0 mg, Strychnos nux-vomica D13 22.0 mg, Sulfur D8 22.0 mg, Ventriculus suis D8 22.0 mg, Veratrum album D4 22.0 mg, Natrium diethylxalacetatum D8 22.0 mg. Excipients: Sodium chloride 19.1 mg, water for injections 1430.0 mg.

Indications: Tablets, injection solution: Disorders of mucous membranes (gastrointestinal, respiratory and genitourinary system and the eye).

Contraindications: Tablets, injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Injection solution:** None.

Side effects: Tablets, injection solution: None have been reported.

Interactions with other medication: Tablets, injection solution: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medication are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. Method of administration: Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Injection solution:** Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅓ of an ampoule 1 to 3x weekly. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅓ of an ampoule daily, and then continue with standard dosage. Method of administration: Mucosa comp.-Heel, Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Tablets, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (52268): Packs containing 50, 100 and 250 tablets. **Injection solution (52546):** Packs containing 5, 10, 50 and 100 ampoules of 2.2 ml each.

Oculoheel® • Eye drops

Compositions: Eye drops: 0.45 g contains: Active ingredients: Chochlearia officinalis D5 110.7 mg, Echinacea D5 110.7 mg, Euphrasia D5 110.7 mg, Pilocarpus D5 110.7 mg. Excipients: Disodium phosphate dihydrate 4.208 mg, Sodium chloride 1.949 mg, Sodium dihydrogen phosphate dihydrate 0.930 mg.

Indications: Eye drops: Irritation and inflammation of the eye, conjunctiva or eyelid.

Contraindications: Eye drops: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Eye drops: The frequent use of eye drops in some patients with severe damage to the cornea may, in isolated cases, lead to corneal calcification.

Side effects: Eye drops: Like all medicinal products, homeopathic medicines may cause side effects. In isolated cases transient local allergic reactions such as redness or burning sensations have been reported. The frequency of these effects is not known.

Interactions with other medication: Eye drops: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Eye drops: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medication are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Eye drops: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Eye drops: Standard Dosage: Adults (and children 12 yrs. and older): 1–2 drops into the affected eye(s) 4x daily, or more often if needed. Pediatric: Below 2 yrs.: 1–2 drops 4x daily as directed [consult a healthcare professional before use]. 2–5 yrs.: 1–2 drops into the affected eye(s) 4x daily, or more often if needed. 6–11 yrs.: 1–2 drops into the affected eye(s) 4x daily, or more often if needed. Method of administration: Instill into the conjunctival sac.

Overdose: Eye drops: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Eye drops (12142): Packages with 15 phials, each containing 0.45 ml of solution.

Important note:

This folder contains helpful health information based on scientific data and is intended for educational purposes only. The information and/or treatment recommendations are not meant as a specific treatment for any individual and should not be construed as a substitute for or a contradiction of professional treatment recommendations by an attending physician or other qualified healthcare professional. Heel is not liable for any damage or loss caused or alleged to be caused, directly or indirectly, based on use of the information provided herein.

Be aware that medication names, indications, and/or formulas may vary from country to country and package inserts may provide country specific information. Therefore, you have to ensure that the content of any material you use is in line with your local law and any other regulatory and/or medical requirements.

Biologische Heilmittel Heel GmbH

Dr.-Reckeweg-Straße 2-4,
76532 Baden-Baden, Germany
www.heel.com

-Heel