

HEEL MEDICATIONS

Women's Health Reference Book



-Heel

Healthcare designed by nature

Main Heel Products For the following symptoms and conditions	Hormeel	Gynäco-heel	Klimakt-Heel	Mulimen	Ovarium comp.	Metro-Adnex-Injeel	Mucosa comp.	Spascu-preeel	Coenzyme comp.	Lympho-myosot	Cerebrum comp.	Echinacea comp.
Disturbances of the female menstrual cycle: amenorrhoea & menorrhagia	Main Product				Main Product				Main Product			
Conditions of the breast: e.g. fibroadenoma and fibrocystic conditions of the breast, mastodynia	Main Product				Main Product					Main Product		
Neuroses, melancholia and depression related to menopause and menopausal transition	Supportive Product	Main Product			Main Product						Main Product	
Premenstrual syndrome & dysmenorrhoea	Supportive Product			Main Product								
Inflammatory conditions of the female genital tract, such as: pelvic inflammatory disease, salpingitis, ovaritis, parametritis, endometritis, myometritis		Main Product			Main Product	Main Product				Main Product		Main Product
Vaginitis of various origins		Main Product				Main Product						
Ovarian cysts		Main Product			Main Product				Main Product			
Physical & autonomic symptoms e.g. hot flushes and palpitations related to menopause and menopausal transition; pruritus vulvae			Main Product		Main Product							
Myoma uteri (uterine fibroids)	Supportive Product											

 Main Product
  Supportive Product

Products and Indications

Product	Indications
Hormeel	<ul style="list-style-type: none"> – Disturbances of the menstrual cycle – Regulation of endocrine functions
Gynäcoheel	<ul style="list-style-type: none"> – Inflammatory conditions of the female genital tract
Klimakt-Heel	<ul style="list-style-type: none"> – Symptoms of the female climacteric syndrome
Mulimen	<ul style="list-style-type: none"> – Premenstrual syndrome and dysmenorrhea
Ovarium compositum	<ul style="list-style-type: none"> – Glandular support in conditions of female hormonal imbalances
Metro-Adnex-Injeel	<ul style="list-style-type: none"> – Inflammatory conditions of the female reproductive organs, dysmenorrhea and mid-cycle pain
Mucosa compositum	<ul style="list-style-type: none"> – Regulation and support of the mucosal membrane functions
Spascupreel	<ul style="list-style-type: none"> – Spasmodic cramps – Spasmodic conditions of the abdomen & urogenital tract
Coenzyme compositum	<ul style="list-style-type: none"> – Stimulation of blocked intracellular enzymatic system
Lymphomyosot	<ul style="list-style-type: none"> – Lymph flow disorders – Recurrent infections – Drainage and Detoxification
Cerebrum compositum	<ul style="list-style-type: none"> – Support in functional and degenerative disorders of the CNS (Central Nervous System)
Echinacea compositum	<ul style="list-style-type: none"> – Immunostimulation – Symptomatic relief in acute infections (especially bacterial)

RESTORES HORMONAL BALANCE

- Disturbances of the menstrual cycle
- Regulation of endocrine functions



Drops

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	10 drops every 1/2-1 hour, Up to 12x daily	10 drops 3x daily

Injection solution


	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	1 ampoule daily	1 ampoule 1-3x weekly

Clinical Evidence

In a multicentric study with 41 physicians and 345 patients the usage, indications, dosages, therapeutic efficacy and tolerance of Hormeel® has been tested.¹

- Hormeel® is an effective treatment of PMS and menopausal syndromes (80 % of the participant reported a very good or good efficacy of PMS and menopausal syndromes)
- Hormeel® has a very good tolerability – 99 % of the participants rated the tolerance as excellent or good
- 87% of patients treated with only Hormeel® achieved very good and good results.

Hormeel® – Clinical Evidence




References

1) Reus, Weiser. Homeopathic Treatment of Gynecological Disorders. Biol Med 1999; 28 (5): 233-236

Clinical Evidence


- **Hormeel®** is an effective treatment as a monotherapy and/or in combination with conventional drug therapy in treating various disturbances of the female menstrual cycle, e.g. PMS, menopausal symptoms, ovarian insufficiency, dysmenorrhea and hormonal dysfunction.¹
- **Hormeel®** is well tolerated by patients both with the oral drops and injection solution.¹
- **Hormeel®** is suitable for long term therapy with minimal side-effects (e.g. treatment time in study included for 1-2 months).¹

Premenstrual Syndrome¹




Very good	27%
Good	58%
Satisfactory	14%
Without success / no data	1%

Menopausal Syndrome¹



Very good	26%
Good	56%
Satisfactory	16%
Without success / no data	2%

Overall tolerance of Hormeel® by patients



References

1) Reus, Weiser. Homeopathic Treatment of Gynecological Disorders. Biol Med 1999; 28 (5): 233-236

Patient Profile

SUZIE, 17-YEAR-OLD | SCHOLAR

(Primary/spasmodic dysmenorrhea)



Medical history: menarche at age 14, no abnormal findings on pelvic examination or imaging studies (i.e. ultrasonography).

Present disorders: Severe spasmodic pains during menstruation that started soon after menarche. The pains last between 48 and 72 hours. The pain is associated with nausea and vomiting. In addition, she feels dizzy and suffers from headaches during this time.

Non-steroidal anti-inflammatories and opioid analgesics do help relieve the pain but leaves her feeling extremely tired. She would like to explore other options rather than a high dose combination oral contraceptive suggested by her family doctor.

RECOMMENDATION

Hormeel®

Mulimen®

Spascupreel® for
spasmodic pain
during the cycle/
menstruation.

Gynäcoheel®

TREATMENT OF INFLAMMATORY CONDITIONS OF THE FEMALE GENITAL TRACT

- Inflammatory conditions of the female genital tract:
Relief of symptoms of female pelvic inflammatory disease (PID), vaginitis, ovaritis



Drops

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	10 drops every 1/2-1 hour, Up to 12x daily	10 drops 3x daily

Patient Profile

INGRID, 24-YEAR-OLD | LEGAL STUDENT

(Vaginitis)



Medical history: just completed a course of antibiotics for a respiratory tract infection.

Present disorders: pruritus of the vagina with accompanying thick, odorless, white discharge as well as marked erythema of the vagina and labia. Pelvic examination and microscopy confirm vulvovaginal candidiasis. Over-the-counter medications have provided only temporary relief.

RECOMMENDATION

Gynäcoheel

Mucosa compositum

Probiotic

Klimakt-Heel®

RELIEF OF FEMALE CLIMACTERIC SYNDROME

- Symptoms due to menopause such as hot flashes, headaches, sweating, fatigue and depressive moods
- A natural alternative or complement to hormone substitution therapy
- Suitable for long-term use
- No risk of dependency or addiction




Tablets

		Acute or initial dosage	Standard dosage
Adults	+ 18	1 tablet every 1/2-1 hour, Up to 12x daily	1 tablet 3x daily

Clinical study

RELIEF OF FEMALE CLIMACTERIC SYNDROME

— Proven effective and well tolerated




In a cohort study of 52 patients with menopausal symptoms, several months of treatment with **Klimakt-Heel®** (1 tablet 3 times a day) produced improvement in symptoms in over 96 percent of cases. No instances of adverse effects were observed.¹

These results were confirmed by an additional study of 45 women whose menopausal symptoms were also treated with **Klimakt-Heel®** tablets. 91 percent of these patients rated the results of therapy as either "very good" or "good".²


Clinical study

RELIEF OF FEMALE CLIMACTERIC SYNDROME

- Klimakt-Heel® leads to pronounced improvement of menopausal symptoms, as measured by Kuppermann scores^{1,2}
- Klimakt-Heel® showed a higher response rate and a better tolerability than a commonly used phytotherapeutic^{1,2}



A) Effects of Klimakt-Heel® compared with Femolene Ultra® treatment on Kupperman score over time of treatment, with measurements taken every four weeks.



B) Cumulative reductions in Kupperman score over the course of the treatment.

References

- 1) Penny, B.L., Razlog, R., Deroukakis, M., Johnston, G. (2009). A comparative study of Klimakt-Heel® and Femolene ultra in the management of typical climacteric symptoms, Health SA Gesundheit 14(1): 144-148. DOI: 10.4102/hsag.v14i1.488
- 2) Penny, B.L. (2005). Therapeutic effects of Klimakt-Heel® compared with a phytotherapeutic preparation for the treatment of menopausal symptoms – a randomized study. J. Biomed. Therapy Fall 2005: 7-10

Patient Profile

KIRSTEN, 55-YEAR-OLD | SECRETARY

married with two children (Menopause)



Medical history: menopause – cessation of menstruation the last 2 years

Present disorders: climacteric syndrome with hot flushes, insomnia and mood swings. She also reports to be feeling more down and despondent since her children recently moved out of home.

Her family doctor suggests the following medicines with bio-regulating properties before prescribing antidepressants and/or hormonal therapy.

RECOMMENDATION

Klimakt-Heel®

Cerebrum
compositum

Mulimen

TREATMENT OF DYSMENORRHEA AND PMS

- Relief of symptoms of dysmenorrhea and of premenstrual syndrome such as mood swings in menstrual disorders



Tablets

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) <div style="float: right; font-size: 2em; font-weight: bold;">+</div> <div style="float: right; border: 1px solid gray; border-radius: 50%; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center;">12</div>	1 tablet every 1/2-1 hour, Up to 12x daily	1 tablet 3x daily
Children 6-11 years	1 tablet every 1-2 hours, Up to 8x daily	1 tablet 2x daily



Drops

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) <div style="float: right; font-size: 2em; font-weight: bold;">+</div> <div style="float: right; border: 1px solid gray; border-radius: 50%; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center;">12</div>	10 drops every 1/2-1 hour, Up to 12x daily	10 drops 3x daily
Children 6-11 years	7 drops every 1/2-1 hour Up to 12x daily	7 drops 3x daily



Injection solution

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) <div style="float: right; font-size: 2em; font-weight: bold;">+</div> <div style="float: right; border: 1px solid gray; border-radius: 50%; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center;">12</div>	1 ampoule daily	1 ampoule 1-3x weekly
Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly

Ovarium compositum

SUPPORT OF THE OVARIAN GLANDS

- Supportive treatment in conditions associated with female hormonal imbalances



Tablets

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) <div style="float: right; border: 1px solid gray; border-radius: 50%; padding: 2px 10px; margin-left: 10px;">+ 12</div>	1 tablet every 1/2-1 hour, Up to 12x daily	1 tablet 3x daily
Children 6-11 years	1 tablet every 1-2 hours, Up to 8x daily	1 tablet 2x daily



Injection solution

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) <div style="float: right; border: 1px solid gray; border-radius: 50%; padding: 2px 10px; margin-left: 10px;">+ 12</div>	1 ampoule daily	1 ampoule 1-3x weekly
Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly

Metro-Adnex-Injeel®

REGULATION OF INFLAMMATORY CONDITIONS OF THE FEMALE REPRODUCTIVE ORGANS

Supportive treatment in:

- Inflammatory conditions of the female reproductive organs
- Dysmenorrhea
- Mid-cycle pain



	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) <div style="float: right; font-size: 2em; font-weight: bold;">+ 12</div>	1 ampoule daily	1 ampoule 1-3x weekly
Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly

Mucosa compositum

FOR HEALTHY FUNCTIONING OF THE MUCOSAL MEMBRANES

- Regulation and support of the mucosal membrane functions



Tablets

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	1 tablet every 1/2-1 hour, Up to 12x daily	1 tablet 3x daily
Children 6-11 years	1 tablet every 1-2 hours, Up to 8x daily	1 tablet 2x daily
Children 2-5 years	1 tablet every 1-2 hours, Up to 6x daily	1 tablet 1-2x daily
Children 0-2 years	1 tablet every 1-2 hours, Up to 4x daily	1 tablet 1x daily



Injection solution



	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	1 ampoule daily	1 ampoule 1-3x weekly
Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly
Children 2-5 years	1/2 ampoule daily	1/2 ampoule 1-3x weekly



Spascupreel®


SOOTHES MENSTRUAL CRAMPS

- Spasmodic cramps
- Spasmodic conditions of the abdomen and urogenital tract



Tablets		Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) 		1 tablet every 1/2-1 hour, Up to 12x daily	1 tablet 3x daily
Children 6-11 years		1 tablet every 1-2 hours, Up to 8x daily	1 tablet 2x daily
Children 2-5 years 		1 tablet every 1-2 hours, Up to 6x daily	1 tablet 1-2x daily
Children 0-2 years		1 tablet every 1-2 hours, Up to 4x daily	1 tablet 1x daily

Injection solution		Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) 		1 ampoule daily	1 ampoule 1-3x weekly
Children 6-11 years 		2/3 ampoule daily	2/3 ampoule 1-3x weekly
Children 2-5 years		1/2 ampoule daily	1/2 ampoule 1-3x weekly

Suppositories		Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) 		1 Suppository every 2 hrs. up to 6x daily	1 Suppository 3x daily
Children 6-11 years		1 Suppository every 2 hrs. up to 4x daily	1 Suppository 2x daily

Coenzyme compositum

SUPPORT OF THE INTRACELLULAR RESPIRATORY CHAIN

- Stimulation of blocked intracellular respiratory enzymatic system



Tablets

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	1 tablet every 1/2-1 hour, Up to 12x daily	1 tablet 3x daily
Children 6-11 years	1 tablet every 1-2 hours, Up to 8x daily	1 tablet 2x daily
Children 2-5 years	1 tablet every 1-2 hours, Up to 6x daily	1 tablet 1-2x daily
Children 0-2 years	1 tablet every 1-2 hours, Up to 4x daily	1 tablet 1x daily



Injection solution




	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	1 ampoule daily	1 ampoule 1-3x weekly
Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly
Children 2-5 years	1/2 ampoule daily	1/2 ampoule 1-3x weekly




Lymphomyosot®




SUPPORT OF THE LYMPHATIC FLOW FOR A FASTER RECOVERY

- Lymph flow disorders
- Recurrent infections
- Drainage and Detoxification



 Tablets		Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12		1 tablet every 1/2-1 hour, Up to 12x daily	1 tablet 3x daily
Children 6-11 years		1 tablet every 1-2 hours, Up to 8x daily	1 tablet 2x daily
Children 2-5 years		1 tablet every 1-2 hours, Up to 6x daily	1 tablet 1-2x daily
Children 0-2 years		1 tablet every 1-2 hours, Up to 4x daily	1 tablet 1x daily

 Drops		Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12		10 drops every 1/2-1 hour, Up to 12x daily	10 drops 3x daily
Children 6-11 years		7 drops every 1/2-1 hour Up to 12x daily	7 drops 3x daily
Children 2-5 years		5 drops every 1/2-1 hour, Up to 12x daily	5 drops 3x daily
Children 0-2 years		3 drops every 1/2-1 hour, Up to 12x daily	3 drops 3x daily

 Injection solution		Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12		1 ampoule daily	1 ampoule 1-3x weekly
Children 6-11 years		2/3 ampoule daily	2/3 ampoule 1-3x weekly
Children 2-5 years		1/2 ampoule daily	1/2 ampoule 1-3x weekly

Cerebrum compositum

SUPPORTIVE TREATMENT OF COGNITIVE FUNCTION

- Support in functional and degenerative disorders of the CNS (Central Nervous System)



Tablets

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	1 tablet every 1/2-1 hour, Up to 12x daily	1 tablet 3x daily
Children 6-11 years	1 tablet every 1-2 hours, Up to 8x daily	1 tablet 2x daily
Children 2-5 years	1 tablet every 1-2 hours, Up to 6x daily	1 tablet 1-2x daily
Children 0-2 years	1 tablet every 1-2 hours, Up to 4x daily	1 tablet 1x daily



Injection solution

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	1 ampoule daily	1 ampoule 1-3x weekly
Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly
Children 2-5 years	1/2 ampoule daily	1/2 ampoule 1-3x weekly

Echinacea compositum

STRENGTHEN YOUR IMMUNITY
IN TIMES OF INFECTION

- Immunostimulation
- Symptomatic relief in acute infections (especially bacterial)



Tablets

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	1 tablet every 1/2-1 hour, Up to 12x daily	1 tablet 3x daily
Children 6-11 years	1 tablet every 1-2 hours, Up to 8x daily	1 tablet 2x daily
Children 2-5 years	1 tablet every 1-2 hours, Up to 6x daily	1 tablet 1-2x daily
Children 0-2 years	1 tablet every 1-2 hours, Up to 4x daily	1 tablet 1x daily



Injection solution

	Acute or initial dosage	Standard dosage
Adults (and children 12 years and older) + 12	1 ampoule daily	1 ampoule 1-3x weekly
Children 6-11 years	2/3 ampoule daily	2/3 ampoule 1-3x weekly
Children 2-5 years	1/2 ampoule daily	1/2 ampoule 1-3x weekly

Product information in alphabetical order

Cerebrum compositum

Cerebrum compositum Summary of Product Characteristics

Tablets • Injection solution

Compositions: **Tablets:** 1 tablet (301.5 mg) contains: Active ingredients: Acidum phosphoricum D10 1 mg, Aconitum napellus D6 1 mg, Aesculus hippocastanum D4 1 mg, Anamirta cocculus D4 1 mg, Arnica montana D28 1 mg, Cerebrum suis D8 1 mg, Cinchona pubescens D4 1 mg, Conium maculatum D12 1 mg, Embryo totalis suis D10 1 mg, Gelsemium sempervirens D6 1 mg, Hepar suis D10 1 mg, Hyoscyamus niger D6 1 mg, Kalium bichromicum D8 1 mg, Kalium phosphoricum D6 1 mg, Magnesium phosphoricum D10 1 mg, Placenta totalis suis D10 1 mg, Ruta graveolens D4 1 mg, Selenium D10 1 mg, Semecarpus anacardium D6 1 mg, Strychnos ignatii D8 1 mg, Sulfur D10 1 mg, Thuja occidentalis D6 1 mg. Excipients: Lactose monohydrate 298.0 mg, Magnesium stearate 1.5 mg. **Injection solution:** 1 ampoule (2.2 g) contains: Active ingredients: Acidum phosphoricum D10 22.0 mg, Aconitum napellus D6 22.0 mg, Aesculus hippocastanum D4 22.0 mg, Ambra grisea D10 22.0 mg, Anamirta cocculus D4 22.0 mg, Arnica montana D28 22.0 mg, Cerebrum suis D8 22.0 mg, Cinchona pubescens D4 22.0 mg, Conium maculatum D4 22.0 mg, Embryo totalis suis D10 22.0 mg, Gelsemium sempervirens D4 22.0 mg, Hepar suis D10 22.0 mg, Hyoscyamus niger D6 22.0 mg, Kalium bichromicum D8 22.0 mg, Kalium phosphoricum D6 22.0 mg, Magnesium phosphoricum D10 22.0 mg, Manganum phosphoricum D8 22.0 mg, Placenta totalis suis D10 22.0 mg, Ruta graveolens D4 22.0 mg, Selenium D10 22.0 mg, Semecarpus anacardium D6 22.0 mg, Strychnos ignatii D8 22.0 mg, Sulfur D10 22.0 mg, Thuja occidentalis D6 22.0 mg. Excipients: Sodium chloride 19.2 mg, water for injections 1672.0 mg.

Indications: **Tablets, injection solution:** Support in functional and degenerative disorders of the CNS (Central Nervous System).

Contraindications: **Tablets, injection solution:** Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: **Tablets:** Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Injection solution:** None. w

Side effects: **Tablets:** None have been reported. **Injection solution:** Allergic (hypersensitivity) reactions (e.g. respiratory distress) may occur in very rare cases (i.e. affects less than 1 in 10,000 users).

Interactions with other medication: **Tablets, injection solution:** No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: **Tablets, injection solution:** For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: **Tablets, injection solution:** No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: **Tablets:** **Standard dosage:** Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. **Method of administration:** Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Injection solution:** **Standard dosage:** Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅔ of an ampoule 1 to 3x weekly. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅔ of an ampoule daily, and then continue with standard dosage.

Method of administration: Cerebrum comp., Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: **Tablets, injection solution:** No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: **Tablets (44180):** Packs containing 50 and 250 tablets.

Injection solution (9475): Packs containing 5, 10, 50 and 100 ampoules of 2.2 ml each.

Coenzyme compositum

Coenzyme comp. Summary of Product Characteristics

Tablets • Injection solution

Compositions: **Tablets:** 1 tablet = 301.5 mg containing: Active ingredients: Acidum cis-aconiticum D8 1.0 mg, Acidum ascorbicum D6 1.0 mg, Acidum citricum D8 1.0 mg, Acidum fumaricum D8 1.0 mg, Acidum alpha-ketoglutaricum D8 1.0 mg, Acidum malicum D8 1.0 mg, Acidum succinicum D8 1.0 mg, Adenosinum triphosphoricum D10 1.0 mg, Beta vulgaris rubra D6 1.0 mg, Coenzym A D8 1.0 mg, Cysteinum D6 1.0 mg, Nadidum D8 1.0 mg, Natrium pyruvicum D8 1.0 mg, Natrium riboflavinum phosphoricum D6 1.0 mg, Nicotinamidum D6 1.0 mg, Pulsatilla pratensis D6 1.0 mg, Pyridoxinum hydrochloricum D6 1.0 mg, Sulfur D10 1.0 mg, Thiaminum hydrochloricum D6 1.0 mg, Acidum thiocticum D6 1.0 mg, Barium oxalsuccinicum D10 1.0 mg, Cerium oxalicum D8 1.0 mg, Hepar sulfuris D10 1.0 mg, Magnesium oroticum dihydricum D6 1.0 mg, Manganum phosphoricum D6 1.0 mg, Natrium diethyloxalacetikum D6 1.0 mg.

Excipients: Lactose monohydrate 293.0 mg, Magnesium stearate 1.5 mg. **Injection solution:** 2.2 g containing: Acidum ascorbicum D6 22.0 mg, Acidum alpha-ketoglutaricum D8 22.0 mg, Acidum cis-aconiticum D8 22.0 mg, Acidum citricum D8 22.0 mg, Acidum fumaricum D8 22.0 mg, Acidum malicum D8 22.0 mg, Acidum succinicum D8 22.0 mg, Adenosinum triphosphoricum D10 22.0 mg, Barium oxalsuccinicum D10 22.0 mg, Beta vulgaris rubra D4 22.0 mg, Coenzym A D8 22.0 mg, Cysteinum D6 22.0 mg, Hepar sulfuris D10 22.0 mg, Nadidum D8 22.0 mg, Natrium pyruvicum D8 22.0 mg, Natrium riboflavinum phosphoricum D6 22.0 mg, Nicotinamidum D6 22.0 mg, Pulsatilla pratensis D6 22.0 mg, Pyridoxinum hydrochloricum D6 22.0 mg, Sulfur D10 22.0 mg, Thiaminum hydrochloricum D6 22.0 mg, Acidum thiocticum D6 22.0 mg, Cerium oxalicum D8 22.0 mg, Magnesium oroticum dihydricum D6 22.0 mg, Manganum phosphoricum D6 22.0 mg, Natrium diethyloxalacetikum D6 22.0 mg. Excipients: Sodium chloride 19.4 mg, water for injections 1628.0 mg.

Indications: Tablets, injection solution: Stimulation of blocked intracellular respiratory enzymatic systems in degenerative diseases.

Contraindications: **Tablets, injection solution:** Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: **Tablets:** Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Injection solution:** None.

Side effects: **Tablets:** None have been reported. **Injection solution:** Allergic (hypersensitivity) reactions (e.g. skin allergies, redness/swelling at the injection site) may occur in very rare cases (i.e. affects less than 1 in 10,000 users).

Interactions with other medication: **Tablets, injection solution:** No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: **Tablets, injection solution:** For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: **Tablets, injection solution:** No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: **Tablets:** **Standard dosage:** Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. **Acute or Initial Dosage:** Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. Method of Administration: Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Injection solution:** **Standard dosage:** Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅔ of an ampoule 1 to 3x weekly. **Acute or Initial Dosage:** Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅔ of an ampoule daily, and then continue with standard dosage.

Method of Administration: Coenzyme comp., Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: **Tablets, injection solution:** No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: **Tablets (41946):** Packs containing 50 and 250 tablets.

Injection solution (8511): Packs containing 5, 10, 50 and 100 ampoules of 2.2 ml each.

Echinacea compositum

Echinacea compositum Summary of Product Characteristics

Tablets • Injection solution

Compositions: Tablets: 301.5 mg contains: Active ingredients: Bryonia D4 0.3 mg, Echinacea D2 0.9 mg, Lachesis D10 0.3 mg, Matricaria recutita D6 0.3 mg, Phytolacca americana D4 0.3 mg, Rhus toxicodendron D4 0.3 mg, Thuja occidentalis D4 0.3 mg, Hepar sulfuris D8 0.3 mg. Excipients: Lactose monohydrate 299.7 mg, Magnesium stearate 1.5 mg. Injection solution: 1 ampoule = 2.2 g contains: Active ingredients: Acidum arsenicosum D8 22 mg, Aconitum napellus D3 22 mg, Argentum nitricum D8 22 mg, Arnica montana D4 22 mg, Baptisia tinctoria D4 22 mg, Bryonia D6 22 mg, Cortisonum aceticum D13 22 mg, Echinacea D5 22 mg, Eupatorium perfoliatum D6 22 mg, Euphorbium D6 22 mg, Gelsemium sempervirens D6 22 mg, Hepar sulfuris D10 22 mg, Hydrargyrum bichloratum D8 22 mg, Lachesis D10 22 mg, Phosphorus D8 22 mg, Phytolacca americana D6 22 mg, Pulsatilla pratensis D8 22 mg, Rhus toxicodendron D4 22 mg, Sanguinaria canadensis D5 22 mg, Sulfur D8 22 mg, Thuja occidentalis D8 22 mg, Zincum metallicum D10 22 mg. Excipients: Sodium chloride 18.5–19.7 mg, Water for injections 1716 mg.

Indications: Tablets, injection solution: Stimulation of the body's intrinsic defense system in cases of inflammations and infections of all tissues and organs.

Contraindications: Tablets, injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. Injection solution: None.

Side effects: Tablets, injection solution: None have been reported.

Interactions with other medication: Tablets, injection solution: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard Dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. **Acute or Initial Dosage:** Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs.: up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. Method of administration: Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. Injection solution: Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅔ of an ampoule 1 to 3x weekly. Acute or Initial Dosage: Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅔ of an ampoule daily, and then continue with standard dosage. Method of administration: Echinacea comp., Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Tablets, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (9820): Packs containing 50 and 250 tablets.

Injection solution (52729): Packs containing 5, 10, 50 and 100 ampoules of 2.2 ml.

Gynäcoheel®

Gynäcoheel® Summary of Product Characteristics

Drops

Compositions: Drops: 100 g contains: Active ingredients: Ammonium bromatum D4 10,0 g, Apis mellifica D4 10,0 g, Aurum iodatum D12 10,0 g, Chamaelirium luteum D4 10,0 g, Liliium lancifolium D4 10,0 g, Melilotus officinalis D3 5,0 g, Palladium metallicum D12 10,0 g, Platinum metallicum D12 10,0 g, Vespa crabro D4 10,0 g, Viburnum opulus D2 5,0 g. Excipients: Water, purified 10,0 g, Contains 35 vol.-% alcohol.

Indications: Drops: Supportive treatment for female pelvic inflammatory disease (PID).

Contraindications: Drops: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Drops: This medicinal product contains 35 vol.-% ethanol (alcohol).

Side effects: Drops: Like all medicinal products, homeopathic medicines can cause side effects in isolated cases, such as transient allergic reactions. The frequency of these effects is not known.

Interactions with other medication: Drops: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Drops: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Drops: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Drops: Standard Dosage: Adults (and children 12 yrs. and older): 10 drops 3x daily. Pediatric: Below 12 yrs.: There is no relevant use of Gynäcoheel, Oral drops in children below 12 years in the indications specified. **Acute or Initial Dosage:** Adults (and children 12 yrs. and older): 10 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage.

Pediatric: Below 12 yrs.: There is no relevant use of Gynäcoheel, Oral drops in children below 12 years in the indications specified.

Method of administration: This medicine should be taken away from meals. For children, add drops to a small amount of water.

Overdose: Drops: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Drops (10010): Drop bottles containing 30 and 100 ml.



Hormeel®

Summary of Product Characteristics

Drops • Injection solution

Compositions: *Drops:* 100 g contains: Active ingredients: Acidum nitricum D4 5,0 g, Aquilegia vulgaris D4 10,0 g, Capsella bursa-pastoris D3 5,0 g, Conyza canadensis D3 5,0 g, Cyclamen purpurascens D4 5,0 g, Cypripedium parviflorum var. pubescens D8 10,0 g, Moschus moschiferus D6 5,0 g, Myristica fragrans D6 10,0 g, Origanum majorana D4 10,0 g, Pulsatilla pratensis D4 5,0 g, Senecio nemorensis ssp. fuchsii D6 5,0 g, Strychnos ignatii D6 5,0 g, Viburnum opulus D3 5,0 g, Calcium carbonicum Hahnemanni D8 10,0 g, Sepia officinalis D6 5,0 g. Excipients: Contains 35 vol.-% alcohol. *Injection solution:* 1 ampoule of 1,1 ml (1,1 g) contains: Active ingredients: Acidum nitricum D4 0,55 mg, Aquilegia vulgaris D4 1,10 mg, Conyza canadensis D3 0,55 mg, Cyclamen purpurascens D4 0,55 mg, Myristica fragrans D6 1,10 mg, Origanum majorana D4 1,10 mg, Pulsatilla pratensis D4 0,55 mg, Strychnos ignatii D6 0,55 mg, Viburnum opulus D3 0,55 mg, Calcium carbonicum Hahnemanni D8 1,10 mg, Sepia officinalis D6 0,55 mg. Excipients: Sodium chloride 10,4 mg, Water for injections 1091,75 mg.

Indications: *Drops, Injection solution:* Disturbances of the menstrual cycle. To regulate endocrine functions.

Contraindications: *Drops, injection solution:* Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: *Drops:* This medicinal product contains 35 vol.-% ethanol (alcohol).

Injection solution: None.

Side effects: *Drops, injection solution:* Like all medicinal products, homeopathic medicines can cause side effects in isolated cases, such as transient allergic reactions. The frequency of these effects is not known.

Interactions with other medication: *Drops, injection solution:* No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: *Drops, injection solution:* For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: *Drops, injection solution:* No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: *Drops: Standard Dosage:* Adults (and children 12 yrs. and older): 10 drops 3x daily. Pediatric: Below 12 yrs.: For children under 12 years please see your doctor. *Acute or Initial Dosage:* Adults (and children 12 yrs. and older): 10 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 12 yrs.: For children under 12 years please see your doctor. **Method of administration:** This medicine should be taken away from meals. For children, add drops to a small amount of water.

Injection solution: Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: Below 12 yrs.: For children under 12 years please see your doctor. *Acute or Initial Dosage:* Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: Below 12 yrs.: For children under 12 years please see your doctor. **Method of administration:** Hormeel, Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: *Drops, injection solution:* No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: *Drops (9774):* Drop bottles containing 30 and 100 ml.

Injection solution (9486): Packs containing 5, 10, 50 and 100 ampoules of 1,1 ml.

Klimakt-Heel®

Klimakt-Heel® Summary of Product Characteristics

Tablets

Compositions: Tablets: 1 tablet contains: Active ingredients: Lachesis D12 90,0 mg, Sepia officinalis D4 30,0 mg, Simarouba cedron D4 30,0 mg, Stannum metallicum D12 60,0 mg, Strychnos ignatii D4 30,0 mg, Sulfur D4 30,0 mg, Sanguinaria canadensis D3 30,0 mg. Excipients: Magnesium stearate 1,5 mg. Contains lactose!

Indications: Tablets: Symptoms of the female climacteric syndrome.

Contraindications: Tablets: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

Side effects: Tablets: Like all medicinal products, homeopathic medicines may cause side effects. In isolated cases transient allergic skin reactions have been reported. The frequency of these effects is not known.

Interactions with other medication: Tablets: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard Dosage: Adults: 1 tablet 3x daily. Pediatric and young adults: Children below 18 yrs. and young adults: There is no relevant use of Klimakt-Heel, Tablets in children below 18 years and young adults in the indications specified.

Acute or Initial Dosage: Adults: 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric and young adults: Children below 18 yrs and young adults: There is no relevant use of Klimakt-Heel, Tablets in children below 18 years and young adults in the indications specified. **Method of administration:** Preferably allow the tablet to dissolve in the mouth, and then swallow. This medicine should be taken away from meals.

Overdose: Tablets: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (9738): Packs containing 50 and 250 tablets.

Lymphomyosot® Summary of Product Characteristics

Tablets • Oral drops • Injection solution

Compositions: **Tablets:** 1 tablet = 301.5 mg containing: Active ingredients: Araneus diadematus D6 15.0 mg, Calcium phosphoricum D12 15.0 mg, Equisetum hiemale D4 15.0 mg, Ferrum iodatum D12 30.0 mg, Fumaria officinalis D4 15.0 mg, Gentiana lutea D5 15.0 mg, Geranium robertianum D4 30.0 mg, Juglans regia ssp. regia D3 15.0 mg, Levothyroxinum D12 15.0 mg, Myosotis arvensis D3 15.0 mg, Nasturtium officinale D4 30.0 mg, Natrium sulfuricum D4 15.0 mg, Pinus sylvestris D4 15.0 mg, Scrophularia nodosa D3 15.0 mg, Smilax D6 15.0 mg, Teucrium scorodonia D3 15.0 mg, Veronica officinalis D3 15.0 mg. Contains lactose. **Oral drops:** 100 g containing: Active ingredients: Araneus diadematus D6 5.0 g, Calcium phosphoricum D12 5.0 g, Equisetum hiemale D4 5.0 g, Ferrum iodatum D12 10.0 g, Fumaria officinalis D4 5.0 g, Gentiana lutea D5 5.0 g, Geranium robertianum D4 10.0 g, Levothyroxinum D12 5.0 g, Myosotis arvensis D3 5.0 g, Nasturtium officinale D4 10.0 g, Natrium sulfuricum D4 5.0 g, Pinus sylvestris D4 5.0 g, Scrophularia nodosa D3 5.0 g, Smilax D6 5.0 g, Teucrium scorodonia D3 5.0 g, Veronica officinalis D3 5.0 g, Excipients: Water, purified 5 g. Contains 35 vol.-% alcohol. **Injection solution:** 1.1 g containing: Active ingredients: Araneus diadematus D6 0.55 mg, Calcium phosphoricum D12 0.55 mg, Equisetum hiemale D4 0.55 mg, Ferrum iodatum D12 1.10 mg, Fumaria officinalis D4 0.55 mg, Gentiana lutea D5 0.55 mg, Geranium robertianum D4 1.10 mg, Levothyroxinum D12 0.55 mg, Myosotis arvensis D3 0.55 mg, Nasturtium officinale D4 1.10 mg, Natrium sulfuricum D4 0.55 mg, Pinus sylvestris D4 0.55 mg, Scrophularia nodosa D3 0.55 mg, Smilax D6 0.55 mg, Teucrium scorodonia D3 0.55 mg, Veronica officinalis D3 0.55 mg. Excipients: Sodium chloride 10.4 mg, water for injections 1089.6 mg.

Indications: Tablets, oral drops, injection solution: Improvement of lymphatic drainage, the non-specific immune defense, and conditions such as benign hypertrophy of lymph nodes, chronic tonsillitis, tonsillar hypertrophy and lymphatic edema.

Contraindications: Tablets, oral drops, injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Oral drops:** This medicinal product contains 35 vol.-% ethanol (alcohol). **Injection solution:** None.

Side effects: Tablets, oral drops, injection solution: Allergic (hypersensitivity) skin reactions may occur in very rare cases (i.e. affects less than 1 in 10,000 users).

Interactions with other medication: Tablets, oral drops, injection solution: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, oral drops, injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, oral drops, injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. **Method of administration:** Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Oral drops: Standard dosage:** Adults (and children 12 yrs. and older): 10 drops 3x daily. Pediatric: Below 2 yrs.: 3 drops 3x daily. 2–5 yrs.: 5 drops 3x daily. 6–11 yrs.: 7 drops 3x daily. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 10 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 3 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. 2–5 yrs.: 5 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. 6–11 yrs.: 7 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage.

Method of administration: This medicine should be taken away from meals. For children, add drops to a small amount of water.

Injection solution: Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅓ of an ampoule 1 to 3x weekly. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅓ of an ampoule daily, and then continue with standard dosage. **Method of administration:** Lymphomyosot, Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Tablets, drops, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (9832): Packs containing 50, 100 and 250 tablets.

Oral drops (10019): Packs containing 30 ml and 100 ml.

Injection solution (9487): Packs containing 5, 10, 50 and 100 ampoules of 1.1 ml each.

Metro- Adnex-Injeel®

Metro-Adnex-Injeel® Summary of Product Characteristics

Injection Solution

Compositions: Injection solution: 1 ampoule of 1,1 ml (=1,1 g) contains: Active ingredients: Apis mellifica D10 1,10 mg, Apis mellifica D30 1,10 mg, Apis mellifica D200 1,10 mg, Cimicifuga racemosa D10 2,20 mg, Cimicifuga racemosa D30 2,20 mg, Cimicifuga racemosa D200 2,20 mg, Hydrargyrum bichloratum D10 1,10 mg, Hydrargyrum bichloratum D30 1,10 mg, Hydrargyrum bichloratum D200 1,10 mg, Lachesis D10 1,10 mg, Lachesis D30 1,10 mg, Lachesis D200 1,10 mg, Lachesis D1000 1,10 mg, Liliium lancifolium D10 1,10 mg, Liliium lancifolium D30 1,10 mg, Liliium lancifolium D200 1,10 mg, Lycopodium clavatum D10 1,10 mg, Lycopodium clavatum D30 1,10 mg, Lycopodium clavatum D200 1,10 mg, Lycopodium clavatum D1000 1,10 mg, Pulsatilla pratensis D10 1,65 mg, Pulsatilla pratensis D30 1,65 mg, Pulsatilla pratensis D200 1,65 mg, Pulsatilla pratensis D1000 1,65 mg, Vespa crabro D10 1,65 mg, Vespa crabro D30 1,65 mg, Vespa crabro D200 1,65 mg. Excipients: Sodium chloride 10,3 mg, Water for injections 1.063,2 mg.

Indications: Injection solution: Inflammatory conditions of the female reproductive organs, dysmenorrhea and mid-cycle pain.

Contraindications: Injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Injection solution: None.

Side effects: Injection solution: Like all medicinal products, homeopathic medicines may cause side effects. In isolated cases transient allergic skin reactions have been reported. The frequency of these effects is not known.

Interactions with other medication: Injection solution: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Injection solution: Standard dosage: Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: Below 6 yrs.: There is no relevant use of Metro-Adnex-Injeel, Solution for injection in children below 6 years in the indications specified. 6-11 yrs.: 2/3 of an ampoule 1 to 3x weekly. **Acute or Initial Dosage:** Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: Below 6 yrs.: There is no relevant use of Metro-Adnex-Injeel, Solution for injection in children below 6 years in the indications specified. 6-11 yrs.: 2/3 of an ampoule daily, and then continue with standard dosage. **Method of administration:** Metro-Adnex-Injeel, Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Injection solution (8554): Packs containing 5, 10, 50 and 100 ampoules of 1,1 ml.

Mucosa compositum

Mucosa compositum Summary of Product Characteristics

Tablets • Injection solution

Compositions: Tablets: 1 tablet = 301.5 mg containing: Active ingredients: Argentum nitricum D6 1.0 mg, Atropa bella-donna D10 1.0 mg, Ceanothus americanus D4 1.0 mg, Hydrastis canadensis D6 1.0 mg, Kalium bichromicum D8 1.0 mg, Kreosotum D12 1.0 mg, Lachesis D10 1.0 mg, Mandragora e radice siccata D10 1.0 mg, Marsdenia cundurango D6 1.0 mg, Momordica balsamina D6 1.0 mg, Mucosa coli suis D8 1.0 mg, Mucosa ductus choledochi suis D8 1.0 mg, Mucosa duodeni suis D8 1.0 mg, Mucosa ilei suis D8 1.0 mg, Mucosa jejuni suis D8 1.0 mg, Mucosa nasalis suis D8 1.0 mg, Mucosa oculi suis D8 1.0 mg, Mucosa oesophagi suis D8 1.0 mg, Mucosa oris suis D8 1.0 mg, Mucosa pulmonis suis D8 1.0 mg, Mucosa pylori suis D8 1.0 mg, Mucosa recti suis D8 1.0 mg, Mucosa vesicae felleae suis D8 1.0 mg, Mucosa vesicae urinariae suis D8 1.0 mg, Oxalis acetosella D6 1.0 mg, Pankreas suis D10 1.0 mg, Phosphorus D8 1.0 mg, Psychotria ipecacuanha D8 1.0 mg, Pulsatilla pratensis D6 1.0 mg, Semecarpus anacardium D6 1.0 mg, Strychnos nux-vomica D13 1.0 mg, Sulfur D8 1.0 mg, Ventriculus suis D8 1.0 mg, Veratrum album D6 1.0 mg. Excipients: Lactose monohydrate 300 mg, Magnesium stearate 1.5 mg.

Injection solution: 2.2 g containing: Active ingredients: Argentum nitricum D6 22.0 mg, Atropa bella-donna D10 22.0 mg, Ceanothus americanus D4 22.0 mg, Hydrastis canadensis D4 22.0 mg, Kalium bichromicum D8 22.0 mg, Kreosotum D10 22.0 mg, Lachesis D10 22.0 mg, Mandragora e radice siccata D10 22.0 mg, Marsdenia cundurango D6 22.0 mg, Momordica balsamina D6 22.0 mg, Mucosa coli suis D8 22.0 mg, Mucosa ductus choledochi suis D8 22.0 mg, Mucosa duodeni suis D8 22.0 mg, Mucosa ilei suis D8 22.0 mg, Mucosa jejuni suis D8 22.0 mg, Mucosa nasalis suis D8 22.0 mg, Mucosa oculi suis D8 22.0 mg, Mucosa oesophagi suis D8 22.0 mg, Mucosa oris suis D8 22.0 mg, Mucosa pulmonis suis D8 22.0 mg, Mucosa pylori suis D8 22.0 mg, Mucosa recti suis D8 22.0 mg, Mucosa vesicae felleae suis D8 22.0 mg, Mucosa vesicae urinariae suis D8 22.0 mg, Oxalis acetosella D6 22.0 mg, Pankreas suis D10 22.0 mg, Phosphorus D8 22.0 mg, Psychotria ipecacuanha D8 22.0 mg, Pulsatilla pratensis D6 22.0 mg, Semecarpus anacardium D6 22.0 mg, Strychnos nux-vomica D13 22.0 mg, Sulfur D8 22.0 mg, Ventriculus suis D8 22.0 mg, Veratrum album D4 22.0 mg, Natrium diethylxalacetikum D8 22.0 mg. Excipients: Sodium chloride 19.1 mg, water for injections 1430.0 mg.

Indications: Tablets, injection solution: Disorders of mucous membranes (gastrointestinal, respiratory and genitourinary system and the eye).

Contraindications: Tablets, injection solution: Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: Tablets: Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. Injection solution: None.

Side effects: Tablets, injection solution: None have been reported.

Interactions with other medication: Tablets, injection solution: No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: Tablets, injection solution: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: Tablets, injection solution: No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: Tablets: Standard dosage: Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 2 yrs.: 1 tablet every 1 to 2 hrs., up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs., up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. **Method of administration:** Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Injection solution: Standard dosage:** Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: 2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅔ of an ampoule 1 to 3x weekly. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅔ of an ampoule daily, and then continue with standard dosage.

Method of administration: Mucosa comp.-Heel, Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Tablets, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: Tablets (52268): Packs containing 50, 100 and 250 tablets.

Injection solution (52546): Packs containing 5, 10, 50 and 100 ampoules of 2.2 ml each.

Mulimen

Mulimen

Summary of Product Characteristics

Tablets • Drops • Injection solution

Compositions: **Tablets:** 1 tablet contains: Active ingredients: Ambra grisea D4 33,33 mg, Calcium carbonicum Hahnemanni D8 33,33 mg, Cimicifuga racemosa D4 33,33 mg, Gelsemium sempervirens D4 33,33 mg, Hypericum perforatum D3 33,33 mg, Sepia officinalis D8 33,33 mg, Urtica urens D3 33,33 mg, Vitex agnus-castus D3 33,33 mg, Kalium carbonicum D4 33,33 mg. Excipients: Magnesium stearate 1,50 mg. Contains lactose! **Drops:** 100 g contains: Active ingredients: Ambra grisea D4 11,11 g, Hypericum perforatum D3 11,11 g, Vitex agnus-castus D3 11,11 g, Cimicifuga racemosa D4 11,11 g, Gelsemium sempervirens D4 11,11 g, Kalium carbonicum D4 11,11 g, Calcium carbonicum Hahnemanni D8 11,11 g, Sepia officinalis D8 11,11 g. Excipients: Ethanol 43 % (m/m) 11,11 g, Contains 54 vol.-% alcohol.

Injection solution: 1 ampoule of 2,0 ml (= 2,0 g) contains: Active ingredients: Cimicifuga racemosa D4 100,00 mg, Gelsemium sempervirens D4 100,00 mg, Hypericum perforatum D3 100,00 mg, Kalium carbonicum D4 100,00 mg, Urtica urens D3 100,00 mg, Ambra grisea D4 100,00 mg, Vitex agnus-castus D3 100,00 mg, Calcium carbonicum Hahnemanni D8 100,00 mg, Sepia officinalis D8 100,00 mg. Excipients: Sodium chloride 5,68 - 6,94 mg, Water for injections 1100,00 mg.

Indications: **Tablets, Drops, Injection solution:** Symptomatic treatment of PMS and dysmenorrhea.

Contraindications: **Tablets, Drops, Injection solution:** Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: **Tablets:** Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Drops:** This medicinal product contains 54 vol.-% ethanol (alcohol). **Injection solution:** None.

Side effects: **Tablets, Drops, Injection solution:** Like all medicinal products, homeopathic medicines can cause side effects in isolated cases, such as transient allergic reactions. The frequency of these effects is not known.

Interactions with other medication: **Tablets, Drops, Injection solution:** No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: **Tablets, Drops, Injection solution:** For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: **Tablets, Injection solution:** No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions. **Drops:** Mulimen, Oral drops have no or negligible influence, at the recommended dosage, on the ability to drive and use machines.

Dosage: **Tablets:** **Standard Dosage:** Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 6 yrs.: There is no relevant use of Mulimen, Oral drops in children below 6 years in the indications specified. 6–11 yrs.: 1 tablet 2x daily. **Acute or Initial Dosage:** Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 6 yrs.: There is no relevant use of Mulimen, Oral drops in children below 6 years in the indications specified. 6–11 yrs.: 1 tablet every 1 to 2 hrs, up to 8x daily, and then continue with standard dosage. **Method of administration:** Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals. **Drops:** **Standard Dosage:** Adults (and children 12 yrs. and older): 10 drops 3x daily. Pediatric: Below 6 yrs.: There is no relevant use of Mulimen, Oral drops in children below 6 years in the indications specified. 6–11 yrs.: 7 drops 3x daily. **Acute or Initial Dosage:** Adults (and children 12 yrs. and older): 10 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 6 yrs.: There is no relevant use of Mulimen, Oral drops in children below 6 years in the indications specified. 6–11 yrs.: 7 drops every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. **Method of administration:** This medicine should be taken away from meals. For children, add drops to a small amount of water. **Injection solution:** **Standard dosage:** Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric Below 6 yrs.: There is no relevant use of Mulimen, Solution for injection in children below 6 years in the indications specified. 6–11 yrs.: ⅓ of an ampoule 1 to 3x weekly. **Acute or Initial Dosage:** Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: Below 6 yrs.: There is no relevant use of Mulimen, Solution for injection in children below 6 years in the indications specified. 6–11 yrs.: ⅓ of an ampoule daily, and then continue with standard dosage. **Method of administration:** Mulimen, Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: **Tablets, injection solution:** No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: **Tablets (9879):** Pack containing 100 tablets.

Drops (9915): Drop bottles containing 50 and 100 ml.

Injection solution (9440): Packs containing 5, 50 and 500 ampoules of 2.0 ml.

Ovarium compositum

Ovarium compositum Summary of Product Characteristics

Tablets • Injection solution

Compositions: **Tablets:** 1 tablet contains: Active ingredients: Acidum cis-aconiticum D10 1,0 mg, Apisinum D8 1,0 mg, Aquilegia vulgaris D4 1,0 mg, Calvatia gigantea D6 1,0 mg, Cypridium parviflorum var. pubescens D6 1,0 mg, Hydrastis canadensis D4 1,0 mg, Hypophysis suis D13 1,0 mg, Kreosotum D8 1,0 mg, Lachesis D10 1,0 mg, Liliium lancifolium D4 1,0 mg, Ovarium suis D8 1,0 mg, Placenta totalis suis D10 1,0 mg, Psychotria ipecacuanha D6 1,0 mg, Pulsatilla pratensis D18 1,0 mg, Tuba uterina suis D10 1,0 mg, Uterus suis D10 1,0 mg, Magnesium phosphoricum D10 1,0 mg, Mercurius solubilis Hahnemanni D10 1,0 mg, Sepia officinalis D10 1,0 mg. Excipients: Lactose monohydrate 297,0 mg, Magnesium stearate 1,5 mg. **Injection solution:** 1 ampoule of 2,2 ml (=2,2 g) contains: Active ingredients: Acidum cis-aconiticum D10 22,0 mg, Apisinum D8 22,0 mg, Aquilegia vulgaris D4 22,0 mg, Calvatia gigantea D6 22,0 mg, Cypridium parviflorum var. pubescens D6 22,0 mg, Hydrastis canadensis D4 22,0 mg, Hypophysis suis D13 22,0 mg, Kreosotum D8 22,0 mg, Lachesis D10 22,0 mg, Liliium lancifolium D4 22,0 mg, Magnesium phosphoricum D10 22,0 mg, Mercurius solubilis Hahnemanni D10 22,0 mg, Ovarium suis D8 22,0 mg, Placenta totalis suis D10 22,0 mg, Psychotria ipecacuanha D6 22,0 mg, Pulsatilla pratensis D18 22,0 mg, Sepia officinalis D10 22,0 mg, Tuba uterina suis D10 22,0 mg, Uterus suis D10 22,0 mg. Excipients: Sodium chloride 19,8 mg, Water for injections 1782,0 mg.

Indications: **Tablets, Injection solution:** Glandular support for conditions of female hormonal imbalance.

Contraindications: **Tablets, injection solution:** Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: **Tablets:** Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Injection solution:** None.

Side effects: **Tablets, injection solution:** Like all medicinal products, homeopathic medicines can cause side effects in isolated cases, such as transient allergic reactions. The frequency of these effects is not known.

Interactions with other medication: **Tablets, injection solution:** No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: **Tablets, injection solution:** For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: **Tablets, injection solution:** No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: **Tablets:** **Standard Dosage:** Adults (and children 12 yrs. and older): 1 tablet 3x daily. Pediatric: Below 6 yrs.: There is no relevant use of Ovarium comp., Tablets in children below 6 years in the indications specified. 6–11 yrs.: 1 tablet 2x daily. Acute or Initial Dosage: Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Pediatric: Below 6 yrs.: There is no relevant use of Ovarium comp., Tablets in children below 6 years in the indications specified. 6–11 yrs.: 1 tablet every 1 to 2 hrs., up to 8x daily, and then continue with standard dosage. **Method of administration:** Preferably allow the tablet to dissolve in the mouth, and then swallow. This medicine should be taken away from meals. **Injection solution:** **Standard dosage:** Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly. Pediatric: Below 6 yrs.: There is no relevant use of Ovarium comp., Solution for injection in children below 6 years in the indications specified. 6–11 yrs.: ⅓ of an ampoule 1 to 3x weekly. **Acute or Initial Dosage:** Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. Pediatric: Below 6 yrs.: There is no relevant use of Ovarium comp., Solution for injection in children below 6 years in the indications specified. 6–11 yrs.: ⅓ of an ampoule daily, and then continue with standard dosage. **Method of administration:** Ovarium comp., Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Overdose: Tablets, injection solution: No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: **Tablets (75527):** Packs containing 50 tablets.

Injection solution (8527): Packs containing 5, 10, 50 and 100 ampoules of 2,2 ml.

Spascupreel®

Spascupreel® Summary of Product Characteristics

Tablets • Injection solution • Suppositories

Compositions: **Tablets:** 1 tablet = 301.5 mg containing: Active ingredients: Aconitum napellus D6 60 mg, Amanita muscaria D4 15 mg, Ammonium bromatum Trit. D4 30 mg, Atropinum sulfuricum D6 30 mg, Citrullus colocynthis D4 30 mg, Cuprum sulfuricum D6 15 mg, Gelsemium sempervirens D6 30 mg, Magnesium phosphoricum D6 30 mg, Matricaria recutita D3 15 mg, Passiflora incarnata D2 15 mg, Veratrum album D6 30 mg; Excipients: Magnesium stearate 1.5 mg; contains lactose. **Injection solution:** 1 ampoule (1.1 g) contains: Active ingredients: Aconitum napellus D6 2.20 mg, Amanita muscaria D4 0.55 mg, Ammonium bromatum D4 1.10 mg, Atropinum sulfuricum D6 1.10 mg, Citrullus colocynthis D4 1.10 mg, Cuprum sulfuricum D6 0.55 mg, Gelsemium sempervirens D6 1.10 mg, Magnesium phosphoricum D6 1.10 mg, Matricaria recutita D3 0.55 mg, Passiflora incarnata D2 0.55 mg, Veratrum album D6 1.10 mg; Excipients: Sodium chloride 10.4 mg, water for injections 1089.0 mg. **Suppositories:** 1 suppository = 2.0 g containing: Active ingredients: Aconitum napellus D5 2.20 mg, Amanita muscaria D3 0.55 mg, Ammonium bromatum D3 1.10 mg, Atropinum sulfuricum D5 1.10 mg, Citrullus colocynthis D3 1.10 mg, Cuprum sulfuricum D5 0.55 mg, Gelsemium sempervirens D5 1.10 mg, Magnesium phosphoricum D6 1.10 mg, Matricaria recutita D2 0.55 mg, Passiflora incarnata D1 0.55 mg, Veratrum album D5 1.10 mg; Excipients: Hard fat 1989.0 mg. **Indications:** Tablets, Injection solution, Suppositories: Spasms of the smooth musculature of the gastrointestinal and the urogenital tract.

Contraindications: **Tablets, Injection solution, Suppositories:** Known allergy (hypersensitivity) to one or more of the ingredients.

Special warnings and special precautions for use: **Tablets:** Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. **Injection solution, Suppositories:** None.

Side effects: **Tablets:** Like all medicinal products, homeopathic medicines may cause side effects. In isolated cases transient allergies (e.g. urticaria, pruritus) as well as nausea have been reported. The frequency of these effects is not known. **Injection solution, Suppositories:** Like all medicinal products, homeopathic medicines can cause side effects in isolated cases, such as transient allergic reactions. The frequency of these effects is not known.

Interactions with other medication: **Tablets, Injection solution, Suppositories:** No interactions have been reported, and none are expected due to the homeopathic dilutions.

Pregnancy and lactation: **Tablets, Injection solution, Suppositories:** For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

Effects on ability to drive and use machines: **Tablets, Injection solution, Suppositories:** No effects on the ability to drive and use machines have been reported, and none are expected due to the homeopathic dilutions.

Dosage: **Tablets:** **Standard dosage:** Adults (and children 12 yrs. and older): 1 tablet 3x daily. Below 2 yrs.: 1 tablet 1x daily. 2–5 yrs.: 1 tablet 1–2x daily. 6–11 yrs.: 1 tablet 2x daily. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage. Below 2 yrs.: 1 tablet every 1 to 2 hrs. up to 4x daily, and then continue with standard dosage. 2–5 yrs.: 1 tablet every 1 to 2 hrs. up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 tablet every 1 to 2 hrs. up to 8x daily, and then continue with standard dosage. **Method of administration:** Preferably allow the tablet to dissolve in the mouth, and then swallow. For children it is possible to crush the tablet and add to a small amount of water. This medicine should be taken away from meals.

Injection solution: **Standard dosage:** Adults (and children 12 yrs. and older): 1 ampoule 1 to 3x weekly.

2–5 yrs.: ½ ampoule 1 to 3x weekly. 6–11 yrs.: ⅔ of an ampoule 1 to 3x weekly. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 1 ampoule daily, and then continue with standard dosage. 2–5 yrs.: ½ ampoule daily, and then continue with standard dosage. 6–11 yrs.: ⅔ of an ampoule daily, and then continue with standard dosage. **Method of administration:** Spascupreel, Solution for injection may be administered by the s.c., i.d., i.m. or i.v. route.

Suppositories: **Standard dosage:** Adults (and children 12 yrs. and older): 1 suppository 3x daily. 6–11 yrs.: 1 suppository 2x daily. **Acute or initial dosage:** Adults (and children 12 yrs. and older): 1 suppository every 2 hrs. up to 6x daily, and then continue with standard dosage. 6–11 yrs.: 1 suppository every 2 hrs. up to 4x daily, and then continue with standard dosage. **Method of administration:** Insert suppository into the rectum.

Overdose: **Tablets, Injection solution, Suppositories:** No cases of overdose have been reported, and none are expected due to the homeopathic dilutions.

Package sizes: **Tablets (9749):** Packs containing 50 and 250 tablets.


Injection solution (8558): packs containing 10 and 100 ampoules of 1.1ml each.

Suppositories (9573): packs containing 12 and 60 suppositories of 2.0 g

REFERENCE BOOKS

Overview

1. Gastrointestinal Disorders
2. Cough and Cold
3. Musculoskeletal Disorders
4. Stress and Sleep
5. Allergies
6. Women's Health



This folder contains helpful health information based on scientific data and is intended for educational purposes only. The information and/or treatment recommendations are not meant as a specific treatment for any individual and should not be construed as a substitute for or a contradiction of professional treatment recommendations by an attending physician or other qualified healthcare professional. Heel is not liable for any damage or loss caused or alleged to be caused, directly or indirectly, based on use of the information provided herein.

Be aware that medication names, indications, and/or formulas may vary from country to country and package inserts may provide country specific information. Therefore, you have to ensure that the content of any material you use is in line with your local law and any other regulatory and/or medical requirements.

Biologische Heilmittel Heel GmbH

Dr. Reckeweg-Straße 2-4
76532 Baden-Baden
Germany

Tel. +49 (0) 722 15 01 00
info@heel.de
www.heel.com

-Heel
Healthcare designed by nature